AD	)		

Award Number: DAMD17-99-1-9059

TITLE: Analysis of Metastasis in Transgenic Mice

PRINCIPAL INVESTIGATOR: Jeffrey Segall, Ph.D.

CONTRACTING ORGANIZATION: Albert Einstein College of Medicine Bronx New York 10461

REPORT DATE: June 2000

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

. REPORT DOCUMENTATION PAGE			OMB No. 074-0188		
Public reporting burden for this collection of informatic the data needed, and completing and reviewing this creducing this burden to Washington Headquarters Se	collection of information. Send comments rvices, Directorate for Information Operation	regarding this burgen estimate of any	other aspect of this col	ection of mornation, including suggestions for	
Management and Budget, Paperwork Reduction Projection 1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 2000	3. REPORT TYPE AND Annual (1 Jun	99 - 31 May 00)		
4. TITLE AND SUBTITLE Analysis of Metastasis i	5. FUNDING NUMBERS DAMD17-99-1-9059				
6. AUTHOR(S) Jeffrey Segall, Ph.D.					
7. PERFORMING ORGANIZATION NAM Albert Einstein College of Medicine Bronx, New York 10461	8. PERFORMING ORGANIZATION REPORT NUMBER				
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSORING / MONITORING	
U.S. Army Medical Research and N Fort Detrick, Maryland 21702-501			AGENCY	REPORT NUMBER	
11. SUPPLEMENTARY NOTES	Report contains	colored photo	s		
12a. DISTRIBUTION / AVAILABILITY S Approved for public release; d		12b. DISTRIBUTION CODE			
13. ABSTRACT (Maximum 200 Words The goal of this project mice. This will make us visualization of the tum the first year of this pexpression construct. W 25) and are in the proceshow GFP fluorescence in carrying the HER2/neu or	is to analyze the e of fluorescently or cells in the pri roject we have gene e have evaluated GF as of testing two commammary ducts and	labeled tumor cel mary tumor as wel erated 3 transgeni TP fluorescence in others. Heterozycare now being cro	ls to allo l as in ta c lines ca n one of th gotes deriv	w in vivo rget organs. For rrying the MMTV-GFP ese lines (founder red from founder 25	
14. SUBJECT TERMS Breast Cancer				15. NUMBER OF PAGES 8 16. PRICE CODE	
				INOL GODE	

19. SECURITY CLASSIFICATION

Unclassified

OF ABSTRACT

18. SECURITY CLASSIFICATION

Unclassified

OF THIS PAGE

OF REPORT

17. SECURITY CLASSIFICATION

Unclassified

20. LIMITATION OF ABSTRACT

Unlimited

Form Approved

### FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

 $\underline{X}$  For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

 $\underline{\text{N/A}}$  In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

 $\underline{\text{N/A}}$  In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI //Signature/

Date

# **Table of Contents**

Cover	1
SF 298	2
Foreword	3
Table of Contents	4
Introduction	5
Body	5-8
Key Research Accomplishments	8
Reportable Outcomes	8
Conclusions	8
References	N/A
Appendices	N/A

### Introduction

The major life threatening aspect of most cancers involves the metastatic spread of cancer. However, methods for dissecting the process of metastasis need further development in order to be able to determine the functions of specific gene products in metastasis. Transgenic mice expressing or lacking specific cancer related genes have provided a powerful method for evaluating mechanisms of tumor formation. This project will develop new techniques for more detailed dissection of the roles of such gene products in the metastatic process, using GFP (green fluorescent protein)-expressing transgenic mice. In the first technical objective, transgenic mice expressing GFP in the mammary gland has been generated. We will first evaluate an expression construct in which the enhanced GFP cDNA with mammalian codon sequences is driven by the full length MMTV promoter and ends with the SV40 intron and polyadenylation sequence. Transgenic mice which have the highest level of GFP expression in the mammary gland and greatest tissue specificity of expression will be chosen for further work. In the second technical objective, these mice will then be used together with image analysis techniques that we have developed in order to dissect the process of metastasis in more detail. By crossing the GFP-mice with mice containing specific mammary tissue-targeted proteins such as HER2/neu, the contributions of specific oncogenes to the metastatic cascade will be determined. The properties of tumor cells at the border of the tumor, the density of tumor cells in the blood, the kinetics of invasion of target organs, and the distribution of individual tumor cells and metastases in target organs such as the lung can be evaluated using such mice. In the third technical objective, the contribution of other proteins to the various stages of metastasis, such as angiogenesis factors, proteases, cytoskeletal proteins, and adhesion proteins will be determined. In this first annual report, we describe nearly completing the first technical objective, and beginning work on the second objective.

## **Body**

The technical objectives of this project are:

<u>Technical Objective 1:</u> To generate transgenic mice carrying constructs targeting GFP to the mammary gland and to evaluate the specificity of GFP expression.

<u>Technical Objective 2:</u> To cross mice expressing GFP in the mammary gland with mice expressing HER2/neu under the control of the MMTV promoter and utilize the GFP expression to evaluate the metastatic properties of the tumors that develop in these mice.

<u>Technical Objective 3:</u> To evaluate the effects of other transgenes or knockouts on the rate limiting step of metastasis of HER2/neu tumors.

The statement of work is:

### STATEMENT OF WORK


- Task 1: To generate transgenic mice expressing green fluorescent protein in the mammary gland (months 1 12).
  - develop expression constructs for mammary targeted expression of GFP (months 1 -3)
  - generate founder mice with the aid of the Transgenic Core Facility (months 3
     -6)

- identify founders that stably transmit the GFP gene, have high mammary specific expression of GFP (months 6 12). This may require testing up to 6 founders with crosses for 2 generations to test transmission and tissue specificity. Up to 240 newborn mice or 72 transgenics may be evaluated.
- Task 2. To cross the GFP transgenic mice with HER2/neu mice and evaluate the metastatic cascade for the tumors that develop (months 12 24).
  - Use at least one test cross to refine assays for analysis of tumor cells in blood and injection of tumor cells for analysis of seeding in lungs (months 13 19). Per cross, if two heterozygote trangenics are crossed, to acquire around 20 double transgenics will require around 80 progeny. 10 single transgenic progeny will also be kept to as controls for background effects, although both transgenics are in the FvB background.
  - perform the cross and allow tumors to develop as tumors reach .5 1 cm, sacrifice mice and analyze the distribution of tumor cells (months 15 - 24).
- Task 3. To cross GFP HER2/neu mice with mice defective in matrilysin or gelsolin and evaluate the contributions of these molecules to the various metastatic steps (months 24 -36).
  - perform the crosses and allow tumors to develop (months 24 30). Per cross, if two heterozygote transgenics are crossed, to acquire around 20 double transgenics will require around 80 progeny. 10 single transgenic progeny will also be kept to as controls for background effects.
  - as tumors reach .5 1 cm, sacrifice mice and analyze the distribution of tumor cells (months 30 36).

# Progress for Year 1


We have nearly fulfilled Task 1 as follows.

 develop expression constructs for mammary targeted expression of GFP (months 1 -3)



We subcloned the coding sequence of the enhanced GFP gene from pEGFPN1 into p206 as follow (Figure 1). EGFP N1 was digested with Not I and p206 was digested with Eco RI. Both digested plasmids were then blunted with Klenow fragment and then digested with HindIII. The eGFPN1 fragment was then ligated into the digested p206, resulting in the construct shown in Figure 1. the MMTV LTR provides specific transcription in the mammary gland while the SV40 splicing and polyadenylation fragment enhances export and translation. The construct was


then digested with SphI and SpeI and the expression construct was used to generate potential founder mice in the Transgenic Mouse Facility at Einstein.



 generate founder mice with the aid of the Transgenic Core Facility (months 3 6)

The transgenic facility provided us with 60 potential founders. We first attempted using PCR to identify transgenic animals but found that more consistent results could be obtained using Southern blotting. Using Southern blotting we identified 4 transgenic mice (Figure 2). Crossing them with nontransgenic Fvb mice, we found that 2 founders (25 and 57) transmitted the transgene efficiently, one transmitted at low efficiency, and 1 has been unable to mate.

• identify founders that stably transmit the GFP gene, have high mammary specific expression of GFP (months 6 - 12). This may require testing up to 6 founders with crosses for 2 generations to test transmission and tissue specificity. Up to 240 newborn mice or 72 transgenics may be evaluated.



Founder 25 has been studied most thoroughly at this point. Analysis of the mammary glands using the fluorescence microscope demonstrates that the ducts and potentially alveolar structures are detectable in transgenic mice (Figure 3) but not in control nontransgenics. In Figure 3, the brightfield image shows a typical mammary gland stained to show ductal and alveolar structures. The GFP fluorescent images on top are from an unstained mammary

gland, showing structures corresponding to the brightfield image. Although we are continuing to test the fluorescence levels of heterozygous animals to determine the relative intensity increase

as well as the fluorescence present in the 57 line, we are beginning crosses with transgenic animals expressing HER2/neu and polyoma middle T antigen in the mammary gland. The polyoma transgenics generate tumors much more rapidly than the HER2/neu animals and will provide us with more opportunities to refine our methods of imaging tumors in transgenic mice.

# **Key Research Accomplishments**

• Generation of transgenic mice expressing GFP in the mammary gland under the control of the MMTV LTR.

# **Reportable Outcomes**

None at this time.

## **Conclusions**

We have fulfilled Task 1 of the project and are on schedule. We have generated transgenic mice expressing GFP in the mammary gland and will next evaluate the opportunity to image tumors using fluorescence microscopy in vivo.